



JANSSEN VACCINE FAST FACTS

NUMBER OF DOSES
ONE (1)



COMMON SIDE EFFECTS

The second dose side effects are more intense and usually start within 1-2 days of the vaccine, and last for 1-2 days.

- Tiredness
- Headache
- Muscle pain
- Chills
- Joint Pain
- Nausea
- Fever
- Feeling unwell
- Swollen lymph nodes
- Injection site pain, swelling, redness

FACT

The vaccine will not give you COVID-19.

A COVID-19 vaccine does not contain any part of the virus and cannot cause an infection or positive result on a viral test.

FACT

Quick development does not mean it's not safe.

Rapid development is the result of decades of scientific progress and collaboration among scientists across the globe.

FACT

It may provide protection against more than one strain.

All viruses develop small mutations over time. Information to date shows that Janssen is effective against multiple strains presently in circulation.

FACT

The vaccine will not change your DNA.

The vaccine is an mRNA vaccine, which does not interact with DNA in any way. It triggers an immune response that produces protective antibodies.

FACT

The vaccine has not been linked to infertility or miscarriage.

There is no evidence that shows the vaccine causes infertility or the likelihood of miscarriage.

FACT

We know exactly what is in the vaccine.

Johnson and Johnson have published the ingredients list for the Janssen vaccine. Speculation that either contains microchips or tracking has been proven false.